

EROS DIFFERENTIAL STUDIES OF CEPHEIDS IN THE MAGELLANIC CLOUDS : STELLAR PULSATION, STELLAR EVOLUTION AND DISTANCE SCALE.

J.P. Beaulieu^{1,2}, D.D. Sasselov^{1,3}

beaulieu@astro.rug.nl, sasselov@cfa.harvard.edu

¹ Institut d'Astrophysique de Paris, CNRS, 98bis Boulevard Arago, F-75014 Paris, France.

² Kapteyn Laboratorium, Postbus 800, 9700 AV Groningen, The Netherlands.

³ Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA.

Abstract

We present a differential study of 500 Magellanic Cepheids with 3 million measurements obtained as a by-product of the EROS microlensing survey. The data-set is unbiased and provides an excellent basis for a differential analysis between LMC and SMC. We investigate the pulsational properties of the Cepheids as a function of metallicity and the constraints they give to stellar pulsation, stellar evolution theory and opacities.

Moreover, our empirical analysis shows a small metallicity effect on the period-luminosity relation which has a significant implication to recent distance estimates based on HST observations of Cepheids in galaxies as far as the Virgo cluster. When the color shift of Cepheids due to metallicity is interpreted as reddening in the determination of the true distance modulus of a target galaxy a systematic error is made. Accounting for that error makes the low- H_0 values (Sandage et al.) higher and the high- H_0 (Freedman et al.) value lower, bringing those discrepant estimates into agreement around $H_0 \approx 70 \text{ kms}^{-1}\text{Mpc}^{-1}$.

1 Introduction

Cepheids are young, intermediate mass (typically $2-10M_\odot$), bright periodic variable stars. These stars have left the main sequence and are in a post core hydrogen-burning and core helium-burning phase. Because of this evolutionary stage, they lie in an area of the HR diagram, the so-called 'instability strip', where their envelopes are unstable to global radial pulsations via the kappa mechanism. Their period of pulsation is tightly correlated with their luminosity, and this period-luminosity relation (PL) is used as a corner stone in deriving local and extragalactic distances for many decades. Most of the recent determinations of the Hubble constant H_0 are based on HST observations of Cepheids as far away as the Virgo cluster and an assumed universal PL relation calibrated in the LMC. Then, secondary distance indicators are used to connect to the large scale smooth Hubble flow.

One of the exciting by-products of the microlensing survey experiments like EROS[40] and MACHO[52, 19] has been the observation in a systematic way of hundreds of thousands stars in the Magellanic Clouds. High quality light curves with excellent phase coverage have been obtained. We present a status report of the analysis of ~ 500 Cepheids in the LMC and in the SMC based on observation performed by EROS.

Magellanic Clouds are a perfect place to study variable stars. We can consider them at the same distance in a given galaxy, and the reddening is small. Therefore the observed differences are intrinsic. Moreover, these two galaxies have different metallicities. We have the opportunity to check whether observations, stellar pulsation and stellar evolution theories give a self-consistent picture for two low metallicities and different chemical content. With the two unbiased samples of Cepheids in the LMC and in the SMC, we have empirically searched for the influence of metallicity on the PL relation.

2 Cepheids as a test of stellar physics, stellar evolution and chemical composition

2.1 EROS observations

Observations were obtained using a 0.4 m f/10 reflecting telescope and a mosaic of 16 buttable CCDs covering an area of 1 degree x 0.4 degree centered in the bar of the LMC. With the first year of

EROS CCD observations (1991-1992) about 2500 images spanning over ~ 130 days, were taken in two broad bandpass filters B_E and R_E centered respectively on 490 and 670 nm. We have obtained 80 000 light curves, with as many as 48 points in a given night. During the 1993-1995 season about 6000 images have been taken of a field centered in the SMC with a pair of very similar filters (B_{E2} , R_{E2}). An accurate transformation between the two systems have been determined[43]. For the 1991-1992 campaign, 9 CCDs were operative, whereas 15 were working for the 1993-1995 campaign.

We searched systematically the EROS database for variable stars using the modified periodogram technique[26, 10], and the AoV method[48, 12]. We delineated the instability strip in the color magnitude diagram to select Cepheids candidates. Then, we performed a visual inspection of the light curves to exclude eclipsing binaries. The remaining 97 stars in the LMC and 450 in the SMC form our sample of Cepheids. One in the LMC ($P_0=3.4438$ days, $P_1 = 2.443$ days) and one in the SMC ($P_0=2.0964$ days, $P_1 = 1.5435$ days) are beat Cepheids. For the SMC sample, improved signal processing techniques will be applied to find more candidates to be compared with the ones discovered by MACHO[2, 52]. We can notice already that the period ratio P_{10} of the double mode found in the SMC is higher than the ones found in the LMC for the same period. It is of special interest to see if the stars that pulsate both in the second and the first overtone modes in the SMC has the same period ratio P_{21} as the LMC ones.

2.2 Classical Cepheids and s-Cepheids.

Since the last decade, Fourier decomposition techniques are proving to be a powerful tool when used for systematic studies of variable stars. They give a quantitative basis for the comparison of non-linear hydrodynamic models and amplitude equation formalism results with observations[15]. A Fourier decomposition of the form $X = X_0 + \sum_{i=1}^N X_i \cos(i\omega_i t + \Phi_i)$ is adopted and the customary quantities $R_{k1} = X_k/X_1$ and $\Phi_{k1} = \Phi_k - k\Phi_1$ are calculated. The amplitude ratio R_{k1} reflects the asymmetry of the variation, and Φ_{k1} the full width at half maximum of the curve. Both R_{k1} and Φ_{k1} are strongly affected by resonances between pulsational modes. For example, it is well known that the so-called Hertzsprung progression of the changing form of Classical Cepheid light curves is due to a 2:1 resonance between the self-excited fundamental mode and the second overtone around ~ 10 days[46, 15]. In the $R_{21} - P$ plane, it corresponds to a minimum at ~ 10 days, and a steep rise followed by $\pi/2$ drop at ~ 10 days in the $\Phi_{21} - P$ plane. Thanks to hydrodynamical models, it has been understood that the center of the resonance corresponds to the tip of the $\Phi_{21} - P$ curve just prior to the steep drop, and at the minimum of the $R_{21} - P$. This has been studied in our Galaxy[46], in the LMC[38], and in the SMC[4]. The resonance center ($P_0 = 10$ days, period ratio $P_{20} = P_2/P_0 = 0.5$) can be used in conjunction with linear pulsation theory to infer the mass and the luminosity of these stars.

For a few decades, Cepheids have been classified in two groups, the Classical Cepheids with their high amplitude, asymmetric and possibly bumpy light curves that follow the Hertzsprung progression, and the s-Cepheids with low amplitude and symmetric light variation. The mapping of the Galactic short period Cepheids[5, 37, 8] shows that s-Cepheids follow different sequences than the classical ones in the $R_{21} - P$ and $\Phi_{21} - P$ planes. The difference has been interpreted as a difference of stellar evolution status (different crossing of the HR diagram), difference in chemical composition, or as consequence of different pulsational mode.

In figure 2 are plotted R_{21} as a function of period ($R_{21} - P$), Φ_{21} as a function of period ($\Phi_{21} - P$) for short period Galactic Cepheids[37, 8], and the EROS LMC[9] and SMC[12] Cepheids.

A morphological separation based on the $R_{21} - P$ plane has been proposed for Galactic Cepheids[5]. We adopted it for LMC and SMC Cepheids, and classify as s-Cepheids the stars lying in the lower part of the diagram. This separation in the $R_{21} - P$ plane is mirrored by a dichotomy in the PL diagram for LMC and SMC Cepheids (see figure 2 for the PL relation of SMC Cepheids). Therefore we conclude that the difference is a consequence of different mode of pulsation, fundamental radial mode (F) for the Classical Cepheid, and first overtone radial mode (1OT) for the s-Cepheids.




Figure 1: This figure gives the $R_{21} - P$, $\Phi_{21} - P$ plane for short period Galactic Cepheids and EROS LMC and SMC Cepheids. S-Cepheids are plotted as filled diamonds, Classical Cepheids are plotted as crosses. Typical error bars are indicated in the upper part of each subfigure.

2.3 s-Cepheids

In the 3 Galaxies, one can notice a rise in the $\Phi_{21} - P$ plane, followed by a sharp drop of $\pi/2$ the so-called 'Z-shape'. This feature is mirrored by a minimum in the $R_{21} - P$ plane. The drop takes place at $P = 3.2 \pm 0.2$ days in our galaxy[37], $P = 2.7 \pm 0.2$ days in the LMC[9] and 2.2 ± 0.2 in the SMC[12]. It has been proposed that this drop is a signature of a 2:1 resonance between the first and the fourth overtone. Up to now, non-linear survey of overtone pulsators have failed to reproduce this feature[6, 7, 1, 45]. One should notice that it is surprising that this resonance could give birth to such a strong feature in Fourier planes because the fourth overtone is strongly damped. Alternative explanations, another class of fundamental mode pulsators with nearly-sinusoidal light curves for period greater than ~ 3 days, tangent bifurcation between first and second overtone pulsations, are not supported by the observed PL relation.

2.4 Classical Cepheids

The Hertzsprung progression and the 2:1 resonance between the second overtone and the fundamental mode for Classical Cepheids is observed in the 3 galaxies. In our Galaxy, it is found to take place in the range 9.5-10.5 days[33]. From MACHO observations[52], it is found to be between 10.5-10.8 days in the LMC. With EROS data, we do not have enough Cepheids to constrain its position in the LMC. From SMC observations, it seems to take place in the range 10.5-11.5 days[12, 4].

Figure 2 show that it is possible to define a common envelope for the Cepheids of the 3 galaxies before the 10 days resonance in the $R_{21} - P$ plane. Can the spread of values in R_{21} show different amount of damping ? Is it correlated with position in the instability strip ? It is also remarkable to notice that the Cepheids of the 3 Galaxies follow the same sequence in the $\Phi_{21} - P$ plane before the resonance. However, Cepheids of shorter period are observed in the LMC than in the galaxy, and even shorter ones are observed in the SMC. Both Cepheids from the 3 Galaxies present similar acoustic properties : the envelopes that can be plotted in the Fourier planes give the limits of pulsational properties of the stars.

The metallicity affects the evolutionary tracks. For high metallicity, the blue loops for low mass stars do not cross the instability strip, whereas they do for low metallicities. With decreasing the metallicity, the lower limit for the mass of a star to become a Cepheid decrease. It gives a strong constraint about the extension of the blue loops in the HR diagrams for different metallicity.

2.5 Discussion about resonance constraints

When going to lower metallicity the ~ 10 days resonance occurs at longer period. For overtone pulsators, the *Z-shape* takes place at shorter period when decreasing metallicity.

From a theoretical point of view, one has to keep in mind that when going to lower metallicities, the evolutionary models will increase the luminosity at fixed mass. This increase of the mass will lead to a diminution of the calculated period ratios. However, when going to lower Z, the opacity bump that drives the pulsation will be smaller, and therefore will increase the period ratios for a fixed mass and luminosity. The final results, the observed position of a resonance center, will be a combination of these two antagonists effects (plus a non-linear shift, which is of the level of a few tenth of a percent in period ratio for the 10 days resonance, for a solar composition and metallicity[33]).

2.6 Using resonance constraints for linear stability envelopes calculations

First of all, we want to stress that up to now, there is no correct non-linear model for Beat Cepheids. Therefore, one cannot know the influence of non-linear shift compared to the results obtained by linear theory. One earlier study[14] has shown that with the current state of art, if one allows a 1 % shift on period ratio due to a non-linear coupling effects, linear stability analysis of hydrostatic envelopes of Beat Cepheids gives weak constraints about mass-luminosity relation. If one plots mass as a function




Figure 2: The Period-Luminosity relation for SMC Cepheids from the EROS database. Classical Cepheids (fundamental mode pulsators) are plotted as crosses, and s-Cepheids (overtone pulsators) are plotted as dots. Two examples of classical (lower right) and s-Cepheids (upper left) light curves are shown in the figure. The same scale for the amplitude has been adopted. The two types of Cepheids can be isolated on morphological basis, and follow different PL relation. In the right figures are plotted the $R_{21} - P$, $\Phi_{21} - P$ for fundamental pulsators in the Galaxy (dots), the LMC (asterix), the SMC (circle). Typical error bars are indicated in the upper part of each subfigure.

of period ratio for a set of models to find the mass and the luminosity of a given beat Cepheid and assume that the non-linear period shift is handled, the nearly horizontal nature of the curves shows that it is difficult to get reliable mass estimates for beat Cepheids. The situation is worse for long period beat Cepheids than for short period ones. However, some efforts have been concentrated on them[33, 47, 18, 52, 34].

The presence of a resonance in the envelope of Cepheids gives an anchor for linear pulsation theory to infer the mass and the radius of these stars. Since the introduction of the new OPAL opacities in 1991, masses determined by Baade-Wesselink analysis, stellar evolution theory and linear pulsation theory are in agreement for the solar metallicity[33, 29]. The new data obtained by EROS and MACHO in the LMC and in the SMC poses a challenge for stellar pulsation and stellar evolution, by inviting to investigate lower metallicities.

One previous survey[14] showed that the implication of the 2:1 resonance for fundamental Cepheids around 10 days and the alleged 2:1 resonance around 3 days for overtone pulsators are difficult to reconcile with the envelope models of low-metallicity ($Z=0.01$ and $Z=0.004$) Cepheids. When going to lower metallicity, the masses derived by our linear calculations are too small. To summarize, a systematic survey of linear stability analysis of Cepheid envelopes have been done[14]. The hydrostatic models are computed with a 200 zones mesh using the OPAL opacity table (OPAL93[28]) and Alexander Ferguson opacities[3] for the chemical composition $X=0.70$ etc $Z=0.02$, $Z=0.01$, $Z=0.004$. We find by iteration the mass (M) and the luminosity (L) that are compatible with a given resonance constraint. We computed the models at different distances from the linear blue edge (0, 200 K and 400K) for the corresponding M and L . To analyses the deficiency of our previous calculations[14], we performed a new survey with the latest OPAL opacities (OPAL95) to test the sensitivity to various effects.

(0) The use of the new OPAL opacities (OPAL95 instead of OPAL93) didn't solve the problems. There was a small increase of masses of a few percent thanks to the inclusion of new metals, and therefore an increase of the driving 'metal bump'. For $Z=0.02$, we have good agreement between the masses




Figure 3: We give the $\log(L/L_{\odot}) - \log(T_{eff})$ diagram for LMC fundamental pulsators (plotted as dots). We derived individual reddening, then transform to standard system and apply a temperature scale[17]. We assume a distance modulus of 18.5 ± 0.13 mag to the LMC[36]. The error bars (1σ) are plotted on a 10.2 day Cepheids. They are dominated by the color equation transformation[27]. One should note that when deriving individual reddening we over-correct for the position in the instability strip. Therefore it appears slightly narrower than it is. The evolutionary tracks are from the Geneva group with $Z=0.008$. The box comes from linear stability analysis of hydrostatic models of resonant Cepheids with period 9.5 and 10.5 days (with $Z=0.01$ and OPAL96 opacities) at 0, 200 K from the linear blue edge. The masses for a 10.5 days Cepheid at the 200 K from the linear blue edge is about $4.9M_{\odot}$ to be compared with the evolutionary track of $4M_{\odot}$ and $5M_{\odot}$.

given by the stellar pulsation theory and the masses given by the evolutionary calculations of the Geneva group[44] and masses determined by Baade Wesselink analysis. To summarize, $Z = 0.03$ with OPAL93 is more or less equivalent to $Z = 0.02$ with OPAL95. However, the discrepancy for LMC and SMC Cepheids mass determinations has not been removed. A gain of ~ 50 % in mass for SMC Cepheids, and ~ 30 % for LMC Cepheids would be needed to be in agreement with the evolutionary calculations.

- (1) We increase the number of zones in our mesh up to 800. The variations we found are smaller than a few tenths of a percent in period ratio.
- (2) We included convection following a classical mixing length theory. Surprisingly, the situation worsen, and we obtained masses even smaller.
- (3) We made a comparison between the two available mixture available with OPAL tables, Grevesse91, and Grevesse93. The main difference is a 5 % increase of the opacity in the range $\log(T) = 5.3 - 6K$ corresponding to the inclusion of more metals, and changes of abundances between the two mixtures, that gets an increase of the mass of about 5 % for a given model. No significant differences have been found between OPAL and OP opacities when using linear pulsation calculation[47]. It would be necessary to get reliable information about the chemical composition of Cepheids in the Magellanic Cloud and in the Galaxy, in order to adjust opacity tables and analyze its influence on the pulsation. Moreover, we have to underline that we use a metallicity of $Z = 0.01$ for LMC (which is a rather high estimate), whereas we use $Z=0.004$ for SMC (which is a rather low estimate).
- (4) We performed an extensive survey with enhanced opacities in different temperature ranges to try to increase the masses of our models. We found that the most sensitive region is (as it could have been expected from inspection of weight functions and our knowledge of stellar pulsation) in the range $\log T = 5 - 5.3$ (metal bump region) where a crude increase of 20 % gives a 30 % increase of the

Table 1: Models computed with OPAL96 and the iterative process described in the text[14, 13]. For each galaxy, we computed a resonant Cepheid of a given period at the blue edge, at the blue edge +200 K, and +400 K. Masses, Luminosities and effective temperatures are given for all the models.

	mass ($M_{\odot}$)	Lum ($L_{\odot}$)	Teff (K)		mass ($M_{\odot}$)	Lum ($L_{\odot}$)	Teff (K)		mass ($M_{\odot}$)	Lum ($L_{\odot}$)	Teff (K)
GAL 10 days	6.06	4243	5648		5.90	3721	5489		5.71	3241	5331
LMC 10.5 days	4.57	4109	5802		4.88	3849	5634		4.96	3464	5469
SMC 11 days	3.63	3882	5873		4.26	3927	5704		4.51	3635	5537

mass for $Z=0.004$. One should note that it is more or less equivalent to increase the metallicity of the models. So it is an 'ad-hoc' or 'cooking' experiment.

(5) In the first survey[14], it has been assumed that the resonance takes place for the 3 galaxies between 9.5-10.5 days, and resonant models for these periods have been computed and discussed. It would be more appropriate to compare a 10 days Cepheid in our Galaxy, with a 10.5 days in the LMC and a 11 days in the SMC. In table 1 are given some models computed by the iterative code. One should keep in mind that the actual fundamental blue edge is shifted to lower temperature by about 100 K, because the first overtone mode is linearly unstable. Far from the blue edge, the results of the calculations are less reliable because of the possible influence of convection that is ignored by the code. At 200 K from the linear blue edge, one can notice that a mass of $5.9 M_{\odot}$ is obtained for the Galaxy whereas, $4.9 M_{\odot}$ is obtained for the SMC, and $4.3 M_{\odot}$ for the SMC.

There is still a problem in reconciling the Cepheid mass determined by our models for the LMC and SMC metallicity using the resonance constraint given by the observations of the ~ 10 days resonance despite our new efforts. A slight increase of the luminosity of the evolutionary calculations, or a small increase of the masses derived by linear stability analysis of hydrostatic models is needed, or an increase of the opacity tables in the 'metal bump region', or the three... We want to underline again that envelope stability analysis are very sensitive to opacities, whereas evolutionary calculations are less. The inclusion of new metals in the opacity tables would increase a bit the metal opacity bump, and therefore may improve a situation, that is already not so bad !

3 Cepheids and the distance scale

3.1 Cepheid distance determination

The long-term *HST* distance scale programs[22, 51, 42] have all adopted the same method[31, 23] based on two-color photometry to determine reddening corrected Cepheid distances. It is assumed that the Cepheid PL relation is universal, and that the Cepheids from the calibrating set and from the target galaxy have the same intrinsic colors. The slopes of a calibrating set of LMC Cepheids are calculated. The PL relation of the Cepheids from the target galaxy is slid against it to derive apparent distance modulus in each band. A true distance modulus of $\mu_{LMC} = 18.5\text{mag}$, a mean reddening of $E(B - V) = 0.10$ and a Galactic extinction law with $R_V = 3.3$ are adopted for the LMC. Then, using the multicolor apparent distance moduli and a Galactic extinction law, the total mean reddening and the true distance modulus of the target galaxy are determined.

However, theory[50, 49] predicts a small metallicity effect on the PL relation as a results of stellar pulsation through the dependence of period on mass and radius; of stellar evolution through the mass luminosity relation; and of stellar atmospheres theory through the relation between effective temperature, absolute magnitude in bandpasses and bolometric correction. As a summary theory predicts that the slopes of the PL relation are independent of Z , only the zero-point is affected, and the metallicity effect depend on band pass. It is predicted to be small, $\delta M_V = -1.1\delta Z$. However,

if one interpretes the color shift due to metallicity as reddening in deriving the true distance modulus of a target galaxy with the method describe above, then one makes a systematic error of $\delta\mu = \delta M_V + R_V \delta(B - V)_0$, $\delta\mu = 28.6\delta Z$, according to theory[50] ($\delta(B - V)_0$ is the color change due to metallicity).

Observational studies were made since the 70s. For example, an intrinsic color shift between LMC and SMC Cepheids has been clearly pointed out[32]. An empirical search for a metallicity effect[20, 25] in three fields of M31 with 36 Cepheids and 152 BVRI measurements have led to ambiguous results : Freedman & Madore[20] claimed that there is no significant effect. Gould[25] reanalyzed their data with a better statistical treatment taking into account the high degree of correlation between the measurements and found an effect. However, due to the number of observations, he did not solve for a wavelength dependent metallicity effect. The current status is that the influence of metallicity is considered to be insignificant[22, 42], or negligible[51], and is ignored in the derivation of extragalactic distances with *HST*.

We used the EROS Cepheids data from LMC and SMC to perform an empirical test for metallicity effect[11, 43]. We have high quality, excellent phase coverage light curves for classical Cepheids and s-Cepheids. Since they pulsate in different modes, they follow different PL relations. In the LMC we keep 51 fundamental pulsators and 27 first overtone pulsators, and 264 fundamental pulsators and 141 first overtone pulsators in the SMC. So, we have two unbiased samples of Cepheids that fill densely the period-luminosity-color (PLC) space, with known difference in metallicity $\delta[Fe/H]_{LMC-SMC} = 0.35$ [30].

3.2 The method

Our method[11, 43] will be applied independently to classical Cepheids and s-Cepheids. First we compute wavelength dependent slopes for LMC and SMC Cepheids. They are the same in both galaxies within the uncertainties.

We search for a metallicity effect that depends upon band pass and model the data in the PLC plane taking into account the high degree of correlation between the measurements[25]. The assumption of our model are constant PL slope with metallicity, and no depth dispersion with the LMC sample. We adopted an LMC mean reddening of $E(B - V) = 0.10$ and a Galactic extinction law with $R_V = 3.3$. The model then has 12 parameters, which are : two linear fits in the PLC space (slopes β_i and b_i , and the zero points α_i and a_i), the distance difference γ_1 , the relative reddening difference γ_2 and a metallicity dependence γ_3^i ($i=1,2$ are the two colours B_E and the R_E). We note that the observed mean magnitude in the i band for a Cepheid p , with period P_p , is $Q_{i,p}$. The residuals for the SMC sample are then :

$$X_{i,p}^3 = Q_{i,p} - (\alpha_i + \beta_i \log P_p + \gamma_1 + \gamma_2 R_i + \gamma_3^i), \quad (1)$$

$$X_{i,p}^4 = Q_{i,p} - [a_i + b_i(Q_{1,p} - Q_{2,p} + \gamma_2(R_2 - R_1) + \gamma_3^1 - \gamma_3^2) + \gamma_1 + \gamma_2 R_i + \gamma_3^i], \quad (2)$$

with $p = 1, \dots, N(n)$ being the number of Cepheids. The form of the covariance matrices of the data, the χ^2 minimization and the iteration procedure are described in [43]. Unlike Gould (1994), we solve for a wavelength dependent metallicity effect and constrain the individual and sample reddenings of the Cepheids by the independently known foreground reddenings of LMC (0.06) and SMC (0.05). Details of the method are given elsewhere[11, 43].

We applied the technique to the Classical and the s-Cepheids independently and obtained the same results. We derived a correction term due to the metallicity dependence to the distance determination method presented in 3.1.

$$\delta\mu = (25.4_{-12}^{+6})[Z - 0.009]$$

This metallicity term is valid in the spectral region covered by EROS filters and can be applied to V, I photometry as shown by the color equation transformation $(V - I) = 1.02(B_E - R_E)$, $\sigma = 0.02\text{mag}$.

3.3 Effect on H_0 determination

Numerous efforts to determine the Hubble constant H_0 have led to dichotomous results, depending on the choice of the cosmological ladder used[24]. However, both the longer distance scale ($H_0 \approx 50 \text{ kms}^{-1}\text{Mpc}^{-1}$), and the shorter distance scale ($H_0 \approx 80 \text{ kms}^{-1}\text{Mpc}^{-1}$) rely strongly on the Cepheid PL relation calibrated in the LMC and the method described in section 3.1.

Using HST VI photometry of Cepheids in M100[22], $H_0 = 80 \pm 17 \text{ km.s}^{-1} \text{ Mpc}^{-1}$ has been derived. M100 Cepheids are metal rich compared to LMC ones. Adopting $[\text{Fe}/\text{H}]=+0.1$ [53], within the uncertainty range of our metallicity effect, this leads to $H_0 = 76 - 70 \text{ km.s}^{-1} \text{ Mpc}^{-1}$. Recently, Cepheid distances to a few more galaxies less metal rich than M100, and even with comparable metallicity to LMC (outer region of M101) have been reported by the same group[23]. It brings down their initial value obtained for M100, and nicely confirms the effect of metallicity on the Cepheid distance scale, since their new distances are less affected by it.

HST VI photometry of Cepheids have been obtained in M96[51] and $H_0 = 69 \pm 8 \text{ km.s}^{-1} \text{ Mpc}^{-1}$. This galaxy has an abundance estimate of $[\text{Fe}/\text{H}] = -0.02$. This imply a small correction in the Cepheid distance that leads to $H_0 \approx 66 \text{ km.s}^{-1} \text{ Mpc}^{-1}$.

Using HST VI photometry of Cepheids in IC4182 and NGC5253[42], $H_0 = 55 \pm 8 \text{ km.s}^{-1} \text{ Mpc}^{-1}$ has been derived. These two galaxies are metal-poor compared to the LMC. We adopt $[\text{Fe}/\text{H}]=-1.3$ [35, 41], caution on the large uncertainties of this metallicity determination, and the correction leads to $H_0 = 59 - 68 \text{ km.s}^{-1} \text{ Mpc}^{-1}$.

The metallicity dependence we found from the LMC/SMC analysis brings all the derivations of H_0 to good agreement around $H_0 \approx 70 \text{ km.s}^{-1} \text{ Mpc}^{-1}$.

It is a pleasure to thank Robert Buchler, Marie-Jo Goupil and Zoltan Koll  th for very pleasant and useful discussions. The modelling part has been done in collaboration with them. We thank Roger Ferlet and Alfred Vidal-Madjar and the members of the EROS team for their continuous support. This work is based on observations held at ESO La Silla.

References

- [1] Albrow M., 1996, this volume
- [2] Alcock C., et al., 1995, *Astron. J.* **109**, 1653
- [3] Alexander D.R., Ferguson J.W., 1994, *Astrophys. J.* **437**, 879
- [4] Andreassen G.K., 1988, *Astr. Astrophys.* **191**, 71
- [5] Antonnello E., Porretti E., 1986, *Astr. Astrophys.* **169**, 149
- [6] Antonnello E., Aikawa T., 1993, *Astr. Astrophys.* **279**, 119
- [7] Antonnello E., Aikawa T., 1995, *Astr. Astrophys.* **302**, 105
- [8] Antonnello E., Morelli P.L., 1996 *Astr. Astrophys.* in press
- [9] Beaulieu J.P., et al., 1995 *Astr. Astrophys.* **303**, 137
- [10] Beaulieu J.P., 1995, in *Astrophysical Application of Stellar Pulsation*, Stobie R.S., Whitelock P.A. (eds.) ASP Conf. Ser. **83**, 260
- [11] Beaulieu J.P., et al., 1996, *Astr. Astrophys.* submitted
- [12] Beaulieu J.P., et al., 1996 *Astr. Astrophys.* in prep

- [13] Beaulieu J.P., et al., 1997 *Astr. Astrophys.* in prep
- [14] Buchler R., et al., 1996, *Astrophys. J.* **462**, L83
- [15] Buchler R., 1996, this volume
- [16] Caldwell J.A.R. & Coulson I.M., 1986, *Mon. Not. R. astr. Soc* **218**, 223
- [17] Chiosi C., Wood P., Capitano N., 1993, *Astrophys. J. Suppl. Ser.*, **86**, 541
- [18] Christensen-Dalsgaard J. & Petersen J.O., 1995, *Astr. Astrophys.* **308**, L661
- [19] Cook K., 1996, this volume
- [20] Freedman W. & Madore B.F., 1990, *Astrophys. J.*, **365**, 186
- [21] Freedman W. & Madore B.F., 1991, *Publ. Astron. Soc. Pacific* **103**, 933
- [22] Freedman W., et al., 1994, *Nature* **371**, 757
- [23] Freedman W., et al. 1996, *STScI Colloquium*, ed. M. Livio, in press
- [24] Fukugita M., Hogan C.J. & Peebles P.J.E. 1993, *Nature* **366**, 309
- [25] Gould A. 1994, *Astrophys. J.* **426**, 542
- [26] Grison P., 1994, *Astr. Astrophys.* **289**, 404
- [27] Grison P., et al., 1995, *Astr. Astrophys. Suppl. Ser.* **109**, 447
- [28] Iglesias C.A., Rogers F.J., Wilson B.G. 1992, *Astrophys. J.* **397**, 717
- [29] Kanbur S. & Simon N.R. 1994, *Astrophys. J.* **420**, 880
- [30] Luck R. & Lambert D. 1992, *Astrophys. J. Suppl.* **79**, 303
- [31] Madore B.F. & Freedman W., 1991, *Publ. Astron. Soc. Pacific* **103**, 933
- [32] Martin N., et al., 1979
- [33] Moskalik P., Buchler R., Marom A., 1992, *Astrophys. J.* **385**, 685
- [34] Morgan S. & Welch D., 1996, *Astron. J.* submitted
- [35] Pagel, B.E.J., Simonson, E.A., Terlevich, R.J., & Edmunds, M.G. 1992, *Mon. Not. R. astr. Soc*, **255**, 325
- [36] Panagia N., et al., 1991, *Astrophys. J.* **380**, L23
- [37] Porretti E., 1994, *Astr. Astrophys.* **285**, 524
- [38] Petersen J.O., & Andreasen G.K., 1987, *Astr. Astrophys.* **180**, 129
- [39] Press, W.H., Teukolsky, S.A., Vetterling, W.T., & Flannery, B.P. 1992, *Numerical Recipes*, 2nd Ed., (Cambridge University Press, Cambridge)
- [40] Renault C., 1996, this volume
- [41] Saha, A., et al., 1996, *STScI Colloquium*, ed. M. Livio, in press
- [42] Sandage A. et al., 1994, *Astrophys. J.* **423**, L13
- [43] Sasselov et al., 1996, *Astr. Astrophys.* submitted
- [44] Schaller G., Schaerer D., Meynet G., Maeder A., 1992, *Astr. Astrophys. Suppl. Ser.* **96**, 269
- [45] Schaller G., Buchler R., & Moskalik P., 1993 unpublished
- [46] Simon N.R. & Lee A.S. 1981, *Astrophys. J.* **248**, 291

- [47] Simon N.R. & Kanbur S. 1994, *Astrophys. J.* **429**, 779
- [48] Schwarzenberg-Czerny, 1989 *Mon. Not. R. astr. Soc* **241**, 153
- [49] Stift, M.J. 1995, *Astr. Astrophys.* **301**, 776
- [50] Stothers N.R. 1988, *Astrophys. J.* **329**, 712
- [51] Tanvir N.R. et al. 1995, *Nature* **377**, 27
- [52] Welch D., 1996, this volume
- [53] Zaritsky, D., Kennicutt, R.C., & Huchra, J. P. 1994, *Astrophys. J.* **420**, 87